

**LUMINESCENT THIN FILMS BASED ON
N-(DIPHENYLPHOSPHORYL)BENZAMIDE Eu^{III} AND Tb^{III} COMPLEXES
FOR LIGHT EMITTING DIODE TECHNOLOGY**

Nataliia Kariaka ^{a*}, Olena Litsis ^a, Yuriy Kolomzarov ^b, Paula Gawryszewska ^c, Sergii Smola ^d,
Nataliia Rusakova ^d, Victor Trush ^a, Tatiana Sliva ^a, Vladimir Amirkhanov ^a

^a*Taras Shevchenko National University of Kyiv, Inorganic Chemistry Department, 12, Lva Tolstogo St., 01601 Kyiv, Ukraine*

^b*V. Lashkaryov Institute of Physics of Semiconductors, NAS of Ukraine, 41 Nauky Ave., 03028 Kyiv, Ukraine*

^c*University of Wrocław, Faculty of Chemistry, 14 F. Joliot-Curie St., 50-383 Wrocław, Poland*

^d*A.V. Bogatsky Physicochemical Institute, NAS of Ukraine, 65080 Odessa, 86 Lustdorfska doroga, Ukraine*

**e-mail: natalia_kariaka@i.ua; phone +38(044)239-3392; fax +38(044)239-3393*

Abstract. A series of lanthanides coordination compounds LnL₃, LnL₃Phen, and LnL₃Dipy with aryl-functionalised carbacylamidophosphate (CAPH) ligand N-(diphenylphosphoryl)benzamide (HL = PhC(O)N(H)P(O)Ph₂) were synthesized for luminescent thin films producing. The bulky organic ligands provide a hydrophobic shell around Ln^{III} ion leading to highly luminescent terbium and europium complexes with long-lived emission. The thin films of the complexes were obtained by spin coating and for the first time by vacuum evaporation. The films exhibit an intensive monochromatic photoemission with decay times equal to 0.78-1.34ms.

Keywords: lanthanide complex, photoluminescence, carbacylamidophosphate, chelate ligand, thin film.