



**CHEMISTRY JOURNAL OF MOLDOVA.**  
*General, Industrial and Ecological Chemistry*

Publication details, including instructions for authors information:  
<http://cjm.asm.md/home>

**SYNTHESIS, STRUCTURE AND INVESTIGATION  
OF GERMANIUM(IV) AND COPPER(II)  
COMPLEXES WITH MALIC ACID AND  
1,10'-PHENANTHROLINE**

Inna Seifullina <sup>a</sup>, Elena Martsinko <sup>a\*</sup>, Elena Chebanenko <sup>a</sup>,  
Eleonora Afanasenko <sup>a</sup>, Viktoriya Dyakonenko <sup>b</sup>, Svitlana Shishkina <sup>b,c</sup>

<sup>a</sup>*I.I. Mechnikov Odessa National University, 65082, Odessa, Ukraine*

<sup>b</sup>*SSI "Institute for Single Crystals", National Academy of Sciences of  
Ukraine, Nauki Ave 60, Kharkiv 61001, Ukraine*

<sup>c</sup>*V.N. Karazin Kharkiv National University, 4 Svobody sq., Kharkiv  
61077, Ukraine*

\**e-mail: lborn@ukr.net*

*Accepted version posted online: 22 May 2017*

**Chemistry Journal of Moldova** is a non-profit and non-commercial scientific journal, which publishes *open access* articles under the [Creative Commons Attribution \(CC-BY\) License](#) that permits use, distribution and reproduction in any medium so long as the original work is properly cited.

**To cite this article:** I. Seifullina, E. Martsinko, E. Chebanenko, E. Afanasenko, V. Dyakonenko, S. Shishkina. Synthesis, Structure and Investigation of Germanium(IV) and Copper(II) Complexes with Malic Acid and 1,10'-Phenanthroline. *Chemistry Journal of Moldova*, 2017, DOI: <http://dx.doi.org/10.19261/cjm.2016.369>

**Disclaimer:** *This is an uncorrected proof version of the manuscript that has been accepted for publication. Chemistry Journal of Moldova provides this version as a service to authors and researchers. Copyediting, typesetting, and the review of the resulting proof will be undertaken on this manuscript before the final publication. During production and pre-press, errors may be found which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.*

## SYNTHESIS, STRUCTURE AND INVESTIGATION OF GERMANIUM(IV) AND COPPER(II) COMPLEXES WITH MALIC ACID AND 1,10'-PHENANTHROLINE

Inna Seifullina<sup>a</sup>, Elena Martsinko<sup>a\*</sup>, Elena Chebanenko<sup>a</sup>, Eleonora Afanasenko<sup>a</sup>,  
Viktoriya Dyakonenko<sup>b</sup>, Svitlana Shishkina<sup>b,c</sup>

<sup>a</sup>I.I. Mechnikov Odessa National University, 65082, Odessa, Ukraine

<sup>b</sup>SSI "Institute for Single Crystals", National Academy of Sciences of Ukraine,  
Nauki Ave 60, Kharkiv 61001, Ukraine

<sup>c</sup>V.N. Karazin Kharkiv National University, 4 Svobody sq., Kharkiv 61077, Ukraine

\*e-mail: lborn@ukr.net

**Abstract.** Two crystalline compounds of germanium(IV) with malic acid (H<sub>3</sub>Mal) and 1,10'-phenanthroline (phen) - [Ge(HMal)<sub>2</sub>(phen)]·phen·2H<sub>2</sub>O (**I**) and [CuCl(phen)<sub>2</sub>][Ge(OH)(HMal)<sub>2</sub>] (**II**) were synthesized for the first time and characterized by elemental analysis, IR spectroscopy and thermogravimetric analysis. Using single crystal X-ray diffraction, two different forms of germanium were elucidated: Ge<sup>4+</sup> (**I**) and hydrolyzed GeOH<sup>3+</sup> (**II**) with a distorted octahedron and pyramid surrounding geometry, respectively.

**Keywords:** germanium(IV) compounds, malic acid, 1,10'-phenanthroline, copper(II) complex, structure.

Received: 22 November 2016/ Revised final: 28 February 2017/ Accepted: 09 March 2017

### Introduction

In the course of our systematical research towards complex formation of germanium(IV) with hydroxycarboxylic acids, which are different by the number and mutual position of hydroxyl and carboxyl groups, our group have previously synthesized and structurally characterized different ligand and heteronuclear coordination compounds with citric [1,2], tartaric [3,4] and xylaric acids [5,6]. Unexpectedly, it was not possible to obtain germanium(IV) complex with the monohydroxydicarboxylic malic acid under the same conditions. At the same time, such research is of great interest because of the properties of malic acid that plays the key role in plants and animals metabolism, and is involved in many biochemical processes, e.g., the Krebs cycle [7]. It is also a versatile polydentate ligand, a large number of mono-, di- and polymeric coordination compounds of different metals with H<sub>3</sub>Mal have been already structurally characterized [8-11].

In the past few years, there were synthesized a number of complex compounds containing the divalent tris(oxalato-O,O')germanate anion [Ge(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>]<sup>2-</sup>, or the neutral bis(oxalato-O,O')germanium fragment [Ge(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>], with cationic transition metal complexes of 1,10-phenanthroline (phen):

[M(phen)<sub>3</sub>][Ge(C<sub>2</sub>O<sub>4</sub>)<sub>3</sub>]·xH<sub>2</sub>O (where M<sup>2+</sup>=Cu<sup>2+</sup>, Fe<sup>2+</sup>, Ni<sup>2+</sup>, Co<sup>2+</sup>), [MGe(phen)<sub>2</sub>(μ<sub>2</sub>-OH)<sub>2</sub>(C<sub>2</sub>O<sub>4</sub>)<sub>2</sub>] (where M<sup>2+</sup>=Cd<sup>2+</sup> and Cu<sup>2+</sup>) [12,13]. Also, there were reported different ligand complexes with 1,10'-phenanthroline and malic acid that link metals to the supramolecular chain: [Zn(HMal)(phen)(H<sub>2</sub>O)], [Co(HMal)(phen)(H<sub>2</sub>O)], [Fe(HMal)(phen)(H<sub>2</sub>O)] [14]. As a result, the idea for obtaining the complexes of germanium(IV) with malic acid by reactions in GeO<sub>2</sub> – H<sub>3</sub>Mal – phen – water and GeO<sub>2</sub> – H<sub>3</sub>Mal – CuCl<sub>2</sub> – phen – water systems has appeared. Here, the synthesis, structural characterization and properties of the new complexes in the solid state and also their comparison with the previously obtained coordination compounds of germanium(IV) with hydroxycarboxylic acids is reported.

### Experimental

The starting reagents for the complexes synthesis were germanium(IV) oxide (GeO<sub>2</sub>, 99.99%), DL-malic acid (H<sub>3</sub>Mal, 99%), 1,10'-phenanthroline (phen, 99%), copper(II) chloride dihydrate (CuCl<sub>2</sub>·2H<sub>2</sub>O, 99.99%) (all ©Sigma Aldrich).

### Instrumentation

Elemental analysis for germanium and copper was performed using inductively coupled

plasma atomic emission spectroscopy (PerkinElmer Optima 2000 DV instrument), chlorine was quantified by mercurimetry. The C, H, and N analysis was performed using Elemental Analyzer CE-440.

*Thermogravimetric analysis* (TGA) was carried out using a Q-1500D device at a heating rate of 10 °C/min, in air, in 20-1000 °C temperature range.

The IR spectra in the 4000-400 cm<sup>-1</sup> range were recorded as potassium bromide pellets on a PerkinElmer Frontier spectrometer (The absorption bands were attributed according to the reference data for the starting malic acid and our earlier obtained germanium(IV) coordination compounds with hydroxycarboxylic acids [1-6] (*s.* – strong, *m.* – middle, *w.* – weak).

*Crystal data* for structures **I** and **II** were measured on an Xcalibur-3 diffractometer (graphite monochromated Mo-K $\alpha$  radiation, CCD detector,  $\varphi$  and  $\omega$ -scanning). The structures were solved by the direct method using SHELXTL package [15]. Full-matrix least-squares refinement against F<sup>2</sup> in anisotropic approximation was used for non-hydrogen atoms. Positions of hydrogen atoms were located from the electron density difference maps and refined by “riding” model with  $U_{iso} = nU_{eq}$  of the carrier atom ( $n = 1.5$  for hydroxyl groups and  $n = 1.2$  for other hydrogen atoms). CCDC 1513407-1513408 contain the supplementary crystallographic data for **I** and **II**, respectively. These data can be obtained, free of charge, via <http://www.ccdc.cam.ac.uk/conts/retrieving.html>, or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or e-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk). Full use of the CCDC package was also made for searching in the CSD Database [16].

### Synthesis

#### [Ge(HMal)<sub>2</sub>(phen)]·phen·2H<sub>2</sub>O (**I**)

Germanium(IV) oxide (26.2 mg, 0.125 mmol), malic acid (33.5 mg, 0.25 mmol) and 1,10'-phenanthroline (45.0 mg, 0.25 mmol) were added to 100 mL of boiling water. Obtained solution was slowly evaporated to 5 mL (80 °C) to give after 24h the result of self-assembly reaction - the solid, from which single crystals suitable for X-ray crystallography were isolated.

Elemental composition, based on single-crystal data for C<sub>32</sub>GeH<sub>28</sub>N<sub>4</sub>O<sub>12</sub> ( $M = 733.17$  g/mol), analytically calculated in %: C 52.37, Ge 9.90, N 7.64, H 3.81; found for the as-synthesized bulk material (in %): C 52.24, Ge 10.05, N 7.18, H 3.72.

TGA data of **I** (weight losses,  $\Delta m$ ): 80-230 °C, endothermic peak 150 °C (-5.0%); 270-360°C, endothermic peak 310°C (-36.7%); 520-690°C, exothermic peak 610°C (-44.0%).

Selected IR data for **I** (in cm<sup>-1</sup>):  $\nu(\text{O-H})=3371\text{m}$ ;  $\nu(\text{C-H ring})=3069\text{w}$ , 2925w;  $\nu(\text{C=O})=1722\text{s}$ ;  $\nu_{as}(\text{COO}^-)=1686\text{s}$ ;  $\nu(\text{C-C, ring})=1588\text{m}$ , 1526m, 1496w;  $\nu_s(\text{COO}^-)=1419\text{m}$ ;  $\nu(\text{C-N})=1343\text{m}$ ;  $\nu(\text{C-O})=1108\text{s}$ , 1057w;  $\delta(\text{C-H})=1041\text{m}$ , 987w, 947w, 905w;  $\nu(\text{Ge-N})=658\text{m}$ ;  $\nu(\text{Ge-O})=717\text{s}$ .

Crystal data for **I** ( $M = 733.17$  g/mol): monoclinic, space group C2,  $a = 17.714(7)$  Å,  $b = 12.830(7)$  Å,  $c = 7.273(3)$  Å,  $\beta = 102.192(5)$ ,  $V = 1615.6(1)$  Å<sup>3</sup>,  $Z = 2$ ,  $T = 294$  K,  $\mu(\text{MoK}\alpha) = 1.019$  mm<sup>-1</sup>,  $D_{calc} = 1.507$  g/cm<sup>3</sup>, 5619 reflections measured, 2898 unique ( $R_{int} = 0.036$ ,  $R_{sigma} = 0.052$ ) which were used in all calculations. The final  $R_1$  was 0.045 ( $I > 2\sigma(I)$ ) and  $wR_2$  was 0.122 (all data).

#### [CuCl(phen)<sub>2</sub>][Ge(OH)(HMal)<sub>2</sub>] (**II**)

In the first step, GeO<sub>2</sub> (104.6 mg, 1 mmol) and malic acid (268.2 mg, 2 mmol) were dissolved in 100 mL of hot water to make first working solution, which was then concentrated to 25 mL (50 °C). In the second step, weighted portions of 1,10'-phenanthroline (360 mg, 2 mmol) and CuCl<sub>2</sub>·2H<sub>2</sub>O (170.5 mg, 1 mmol) were added to 25 mL of ethanol and then heated at 40 °C until the reagents were completely dissolved. In the third step, two working solutions cooled to the room temperature were mixed to allow obtaining the mild after 2 days, from which blue crystals were collected mechanically and analyzed by X-ray crystallography.

Elemental composition, based on single-crystal data for C<sub>32</sub>ClCuGeH<sub>25</sub>N<sub>4</sub>O<sub>11</sub> ( $M = 813.14$ ) analytically calculated in %: C 47.23, Cl 4.36, Cu 7.80, Ge 8.93, H 3.07, N 6.88; found for the as-synthesized bulk material (in %): C 47.95, Cl 4.10, Cu 8.05, Ge 8.47, H 3.14, N 7.12.

TGA data of **II** (weight losses,  $\Delta m$ ): 150-220°C, endothermic peak 170°C (-2.5%); 260-340°C, endothermic peak 300°C (-44.1%); 400-570°C, exothermic peak 550°C (-10.0%); 600-880°C, exothermic peak 710°C (-20.8%).

Selected IR data for **II** (in cm<sup>-1</sup>):  $\nu(\text{C-H, aromatic})=2919\text{w}$ ;  $\nu(\text{C=O})=1706\text{m}$ ;  $\nu_{as}(\text{COO}^-)=1650\text{s}$ ;  $\nu(\text{C-C, ring})=1587\text{w}$ , 1519m;  $\nu_s(\text{COO}^-)=1428\text{m}$ ;  $\nu(\text{C-N})=1341\text{m}$ ;  $\nu(\text{C-O})=1170\text{m}$ , 1106w;  $\delta(\text{C-H})=961\text{w}$ , 907w;  $\delta(\text{Ge-O-H})=853\text{m}$ ;  $\nu(\text{Ge-O})=668\text{m}$ .

Crystal data for **II** ( $M = 813.14$  g/mol): monoclinic, space group P2<sub>1</sub>,  $a = 12.413(3)$  Å,  $b = 9.823(3)$  Å,  $c = 26.538(4)$  Å,  $\beta = 103.105(6)$ ,

$V=3151.5(3) \text{ \AA}^3$ ,  $Z=4$ ,  $T=294 \text{ K}$ ,  $\mu(\text{MoK}\alpha)=1.784 \text{ mm}^{-1}$ ,  $D_{\text{calc}}=1.714 \text{ g/cm}^3$ , 26405 reflections measured, 10652 unique ( $R_{\text{int}}=0.080$ ,  $R_{\text{sigma}}=0.099$ ), which were used in all calculations. The final  $R_1$  was 0.060 ( $I > 2\sigma(I)$ ) and  $wR_2$  was 0.130 (all data).

## Results and discussion

Complexes **I** and **II** are crystalline solids with the molar ratio Ge : malate : phen = 1:2:2 (**I**) and Ge : malate : Cu : phen = 1:2:1:2 (**II**) that are stable in air.

The thermal decomposition of **I** starts with an endotherm peak in the temperature range 80-230 °C, which is coming along the elimination to the gas phase of two molecules of crystal water (calc.  $\Delta m=4.92\%$ ). While heating, further in a temperature range from 270 to 360 °C, complex **I** eliminates one molecule of phenanthroline and two  $\text{CO}_2$  molecules from the malate ligands (calc.  $\Delta m=36.56\%$ ).

According to the calculated weight loss of the thermogravimetric curve of complex **II**, the first endotherm peak is accompanied by elimination of one water molecule to the gas phase (calc.  $\Delta m=2.2\%$ ). That molecule is formed by an interaction of the carboxyl proton with the inner sphere OH group bound to germanium [3]. The second endotherm peak from the thermogravimetric curve of **II** corresponds to the loss of two phenanthroline molecules (calc.  $\Delta m=44.3\%$ ).

Finally, complexes **I** and **II** undergo oxidative thermal degradation and combustion of the organic part of their molecules, with the corresponding one (for **I**) and two exothermic peaks (for **II**).

The presence in IR spectra of **I** and **II**  $\nu(\text{C}=\text{O})$ ,  $\nu_{\text{as}}(\text{COO}^-)$  and  $\nu_{\text{s}}(\text{COO}^-)$  absorption bands, which are typical for  $\text{COOH}$  and  $\text{COO}^-$  groups, shows the presence of nonequivalent carboxyl groups (free and bound) in the complexes. The conclusion about how these groups are bonded with germanium is also made on the basis of the band corresponding to the Ge–O stretching vibrations, which is emergence in the IR spectra of complexes. The presence of the band responsible for the Ge–O–H bending vibrations implies the presence of the hydrolyzed form of germanium in the complex **II**.

The absorption band at  $\sim 1340 \text{ cm}^{-1}$  can be assigned to the  $\nu(\text{CN})$  heterocycle, the bands at 1588, 1526, 1496  $\text{cm}^{-1}$  (for **I**) and 1587, 1519 (for **II**) - to the  $\nu(\text{C}-\text{C})$  phenanthroline ring vibrations. The deformation vibrations  $\delta(\text{C}-\text{H})$  of the aromatic rings: planar vibrations at  $1041 \text{ cm}^{-1}$ ;

non-planar vibrations at  $987-905 \text{ cm}^{-1}$  region were found [17,18].

In the structures of complexes **I** and **II** two different forms of germanium are implemented:  $\text{Ge}^{4+}$  (**I**) and hydrolyzed  $\text{GeOH}^{3+}$  (**II**), wherein neutral complex and anion are formed, respectively. In **I** coordination number is 6, in **II** – 5, where germanium atom is coordinated with two malate anions  $\text{HMal}^{2-}$  in both cases and binds with 1,10<sup>7</sup>-phenanthroline in **I** and with hydroxyl group in **II**.

The compound **I** exists as a monohydrate of co-crystal with 1,10<sup>7</sup>-phenanthroline in 1:1 ratio (Figure 1). Molecules of complex and uncoordinated 1,10<sup>7</sup>-phenanthroline are in particular position on the axis of second-order symmetry.




Figure 1. Molecular structure of **I**.

The coordination polyhedron of Ge is a distorted octahedron. The Ge–O bond lengths are 1.818(3) Å and 1.912(5) Å, the Ge–N bond length is 2.057(3) Å. The O–Ge–O, O–Ge–N, N–Ge–N valence angles vary within  $79.8(4)^\circ \div 94.9(2)^\circ$  (Table1).

In crystal structure of the molecules of coordination compound **I** and water are connected with intermolecular hydrogen bonds (Table 2) and form the three-dimensional network (Figure 2).

Table 1  
Bond length (Å) and bond angles (°) in structures **I** and **II** (<sup>1</sup>-x,1-y,-z).

Structure I	
Ge1-O1	1.818(3)
Ge1-O2	1.911(5)
Ge1-N1	2.057(6)
O1 <sup>1</sup> -Ge1-O2	91.6(2)
O1-Ge1-O2	87.1(2)
O1-Ge1-N1 <sup>1</sup>	94.9(2)
O1-Ge1-N1	86.5(2)
O2-Ge1-O2 <sup>1</sup>	93.6(3)
O2-Ge1-N1 <sup>1</sup>	93.7(9)

<sup>1</sup>1-x, +y, 1-z

Continuation of Table 1

Structure II		
	Molecule A	Molecule B
Cu1-Cl1	2.315(3)	2.315(3)
Cu1-N1	2.103(7)	1.986(8)
Cu1-N2	1.994(8)	2.103(8)
Cu1-N3	2.107(8)	1.990(9)
Cu1-N4	1.987(8)	2.105(7)
Ge1-O1	1.793(7)	1.781(6)
Ge1-O2	1.910(6)	1.900(6)
Ge1-O6	1.773(7)	1.779(6)
Ge1-O7	1.936(6)	1.937(6)
Ge1-O11	1.741(7)	1.744(7)
N1-Cu1-Cl1	125.4(2)	123.6(2)
N1-Cu1-N3	112.5(3)	114.5(3)
N2-Cu1-Cl1	95.2(3)	91.8(3)
N2-Cu1-N1	80.6(3)	80.9(3)
N2-Cu1-N3	94.1(3)	97.9(3)
N3-Cu1-Cl1	122.0(2)	121.8(2)
N4-Cu1-Cl1	92.1(2)	92.8(3)
N4-Cu1-N1	96.1(3)	95.0(3)
N4-Cu1-N3	81.0(3)	81.1(3)
O1-Ge1-O2	87.7(3)	87.7(3)
O1-Ge1-O7	88.0(3)	86.7(3)
O6-Ge1-O1	124.4(3)	124.8(3)
O6-Ge1-O2	88.9(3)	90.8(3)
O6-Ge1-O7	86.9(3)	86.9(3)
O11-Ge1-O1	117.9(3)	119.5(3)
O11-Ge1-O2	98.1(3)	98.2(3)
O11-Ge1-O6	117.5(3)	115.3(3)
O11-Ge1-O7	91.0(3)	90.4(3)

Table 2

## Geometric parameters of the hydrogen bonds in structures I and II.

D-H...A	d(H-A) (Å)	d(D-A) (Å)	D-H-A (°)
Structure I			
O5-H5...O6	1.77	2.586(8)	170.2
O6-H6B...O3 <sup>1</sup>	2.01	2.839(10)	165.3
Structure II			
O5A-H5A...O8A <sup>2</sup>	1.98	2.695(11)	144.7
O10A-H10D...O4A	1.96	2.743(15)	160.1
O11A-H11D...O3A <sup>3</sup>	1.98	2.783(10)	142.6
O5B-H5B...O8B <sup>4</sup>	1.87	2.667(10)	163.6
O10B-H10C...O4B	2.18	2.974(15)	162.4
O11B-H11C...O3B <sup>5</sup>	2.24	2.840(10)	141.6
<sup>1</sup> 1/2-x, 1/2+y, 1-z; <sup>2</sup> 1-x, -1/2+y, 1-z; <sup>3</sup> -x, 1/2+y, 1-z; <sup>4</sup> 1-x, 1/2+y, 2-z; <sup>5</sup> -x, -1/2+y, 2-z			

Uncoordinated molecules of 1,10'-phenanthroline occupy cavities between neighborhood complexes. The distance between the  $\pi$ -systems of uncoordinated and coordinated phenanthroline molecules (3.38 Å) allows presuming the stacking interactions existence.

The negative charge of complex anion  $[\text{GeOH}(\text{HMal})_2]^-$  (Figure 3) in structure **II** is compensated by the complex cation  $[\text{CuCl}(\text{phen})_2]^+$  (Figure 4). In the independent part of unit cell, cations and anions of two types were found: A and B with several geometrical parameters difference.




Figure 2. Molecules package in I.

The coordination polyhedron of germanium atom in anion is a pyramid with distorted square in the base. The base of the pyramid is formed by oxygen atoms of deprotonated hydroxyl and carboxyl groups of malic acid (pseudo-torsion angle  $\text{O}(1)\dots\text{O}(2)\dots\text{O}(6)\dots\text{O}(7)$  shows the distortion rate of the base equals to  $30^\circ$  in anions A and B). There is the hydroxyl anion at the top of the pyramid. The negative charge of germanium coordination sphere leads to the changes in bond lengths of Ge-O (they vary within  $1.741(7)\div 1.935(7)$  Å in anion A and  $1.744(7)\div 1.937(6)$  Å in anion B (Table 1)). Valence angles of O-Ge-O bond have values within  $86.9(3)\div 117.9(3)^\circ$  in anions A and B (ideal value is  $90^\circ$ ). The position of the vacant carboxyl groups in anion is stabilized with an intermolecular hydrogen bond  $\text{O}(10)\text{-H}\dots\text{O}(4)$  ( $\text{H}\dots\text{O}$  1.96 Å  $\text{O-H}\dots\text{O}$   $160^\circ$  in anion A and  $\text{H}\dots\text{O}$  2.18 Å  $\text{O-H}\dots\text{O}$   $162^\circ$  in anion B).

The coordination polyhedron of Cu atom in complex cation of **II** represents the distorted trigonal bipyramid (Figure 4).

The N(1), N(3), Cl(1) atoms in cation A or N(2), N(4), Cl(1) in cation B are located in the equatorial planes, the N(2) and N(4) atoms in A or N(1) and N(3) in B, respectively, are located in the axial positions. The Cu-N bond lengths change within  $1.988(8)\div 2.106(9)$  Å in A and  $1.987(8)\div 2.104(8)$  Å in B, at the same time Cu-Cl length is equal to  $2.315(3)$  Å in cation A and  $2.314(3)$  Å in cation B (Table 1).






Figure 3. Molecular structure of anion  $[\text{GeOH}(\text{HMal})_2]^-$  in **II**.




Figure 4. Molecular structure of cation  $[\text{CuCl}(\text{phen})_2]^+$  in **II**.




Figure 5. Crystal packing of compound **II** (projections along the *b* crystallographic axis).

The valence angles in cation A vary within  $112.6(3)\text{--}125.4(2)^\circ$  in the equatorial direction (the ideal value is  $120^\circ$ ) and within  $80.6(3)\text{--}96.1(3)^\circ$  in the axial direction (the ideal value is  $90^\circ$ ), in cation B  $114.5(3)\text{--}123.7(2)^\circ$  in the equatorial direction and within  $80.9(3)\text{--}97.9(3)^\circ$  in the axial direction.

In the crystal structure of **II** anions and cations form three types of alternating layers, parallel along the *ab* crystallographic plane (Figure 5): the first layer type is formed by anions A, the second layer type includes both cations (A and B) and the third layer type is organized by anions B. All anions within the layers are connected by the O-H...O intermolecular hydrogen bonds (Table 2).

## Conclusions

Using 1,10'-phenanthroline, for the first time it was possible to get two different types of malatogermanate complexes of germanium: the different-ligand compound **I** by the self-assembly method from the water solution and the different-metal compound **II** by stepwise synthesis method from the water-ethylene solution. The feature of compound **I** is that 1,10'-phenanthroline enters the inner sphere and coordinates in a bidentate fashion to Ge(IV) by two N atoms, though in water solutions Ge rather coordinates with oxygen-containing organic molecules, such as malic acid in complex anion  $[\text{Ge}(\text{OH})(\text{HMal})_2]^-$  (**II**). The big interest also represents the differences in crystal structures of synthesized

complexes: three-dimensional net, in the cavities of which uncoordinated 1,10'-phenanthroline molecules are located in **I**, and fiber structure of alternating topological isomeric cations and anions A and B in **II**.

## References

- Seifullina, I.I.; Pesaroglo, A.G.; Minacheva, L.Kh.; Martsinko, E.E.; Sergienko, V.S. Bis(citrato)germanate Complexes with Organic Cations: Crystal Structure of  $(\text{HNic})_2[\text{Ge}(\text{HCit})_2] \cdot 3\text{H}_2\text{O}$ . Russian Journal of Inorganic Chemistry, 2006, 51, pp. 1892–1899. DOI: [10.1134/S0036023606120096](http://dx.doi.org/10.1134/S0036023606120096).
- Seifullina, I.I.; Ilyukhin, A.B.; Martsinko, E.E.; Sergienko, V.S.; Chebanenko, E.A. Products of reaction between bis(citrate)hydroxogermanic acid and organic molecules. Molecular and crystal structure of  $(\text{HNad})_2[\text{Ge}(\text{HCit})_2] \cdot 4\text{H}_2\text{O}$ . Russian Journal of Inorganic Chemistry, 2015, 60, pp. 33–37. DOI: [10.1134/S0036023615010143](http://dx.doi.org/10.1134/S0036023615010143).
- Minacheva, L.Kh.; Seifullina, I.I.; Ilyukhin, A.B.; Martsinko, E.E.; Sergienko, V.S.; Chebanenko, E.A.; Churakov, A.V. Strategy for the synthesis of di- and polymer tartratogermanates with single-charge cations. Crystal structures of  $\text{K}_2[\text{Ge}_2(\text{OH})_2(\mu\text{-Tart})_2] \cdot 4.5\text{H}_2\text{O}$  and  $(\text{NH}_4)_{2n}[\text{Ge}_2(\mu\text{-O})(\mu\text{-Tart})_2]_n \cdot n\text{MeCN} \cdot n\text{H}_2\text{O}$ . Russian Journal Coordination Chemistry, 2013, 39, pp. 751–757. DOI: [10.1134/S1070328413110043](http://dx.doi.org/10.1134/S1070328413110043).
- Martsinko, E.E.; Seifullina, I.I.; Minacheva, L.Kh.; Pesaroglo, A.G.; Sergienko, V.S. Synthesis, properties, and molecular and crystal structure of diantipyrylmethanium bis( $\mu$ -tartrato)dihydroxy digermanate(IV) tetrahydrate  $(\text{HDam})_2[\text{Ge}_2(\mu\text{-L})_2(\text{OH})_2] \cdot 4\text{H}_2\text{O}$ . Russian Journal of Inorganic Chemistry, 2008, 53, pp. 1694–1702. DOI: [10.1134/S0036023608110053](http://dx.doi.org/10.1134/S0036023608110053).
- Martsinko, E.E.; Minacheva, L.Kh.; Seifullina, I.I.; Chebanenko, E.A.; Sergienko, V.S.; Churakov, A.V. Synthesis and characterization of cobalt(II) and manganese(II) xylyratogermanates: the molecular and crystal structures of the  $[\text{M}(\text{H}_2\text{O})_6][\text{Ge}_2(\mu\text{-L})_2\{\text{M}(\text{H}_2\text{O})_2\}_2] \cdot 4\text{H}_2\text{O} \cdot n\text{CH}_3\text{CN}$  Complexes ( $\text{M}=\text{Co}$ ,  $n=0$ ;  $\text{M}=\text{Mn}$ ,  $n=1$ ). Russian Journal of Inorganic Chemistry, 2013, 58, pp. 152–159. DOI: [10.1134/S0036023613020174](http://dx.doi.org/10.1134/S0036023613020174).
- Martsinko, E.E.; Minacheva, L.Kh.; Seifullina, I.I.; Pesaroglo, A.G.; Sergienko, V.S. Heteronuclear alkali metal bis( $\mu$ -trihydroxoglutarato)dihydroxy digermanates(IV). The crystal and molecular structure of  $\text{K}_4[\text{Ge}_2(\mu\text{-Thgl})_2(\text{OH})_2] \cdot 4\text{H}_2\text{O}$ . Russian Journal of Inorganic Chemistry, 2012, 57, pp. 343–349. DOI: [10.1134/S0036023612030199](http://dx.doi.org/10.1134/S0036023612030199).
- Sigel, H.; Sigel, A.E. Eds. Metal ions in biological systems: Vanadium and its role in life. M. Dekker: New York, 1995, pp. 147–209.
- Biagioli, M.; Strinna-Erre, L.; Micera, G. Molecular structure, characterization and reactivity of dioxo complexes formed by vanadium(V) with  $\alpha$ -hydroxycarboxylate ligands. Inorganica Chimica Acta, 2000, 310, pp. 1–9. [http://dx.doi.org/10.1016/S0020-1693\(00\)00263-2](http://dx.doi.org/10.1016/S0020-1693(00)00263-2).
- Xie, F.-T.; Duan, L.-M.; Xu, J.-Q.; Ye, L.; Liu, Y.-B.; Hu, X.-X.; Song, J.-F. Solvothermal syntheses and structural characterisation of three isostructural 3D metal-malate coordination polymers:  $\{[\text{M}(\text{C}_4\text{H}_4\text{O}_5)(\text{H}_2\text{O})] \cdot \text{H}_2\text{O}\}_n$  ( $\text{M}=\text{Co}^{\text{II}}$ ,  $\text{Ni}^{\text{II}}$ ,  $\text{Co}^{\text{II}}/\text{Ni}^{\text{II}}$ ). European Journal Inorganic Chemistry, 2004, 22, pp. 4375–4379. DOI: [10.1002/ejic.200400556](http://dx.doi.org/10.1002/ejic.200400556).
- Zhou, Z.-H.; Ye, J.-J.; Deng, Y.F.; Wang G.; Gao, J.-X.; Wan, H.-L. Monomeric and polymeric nickel complexes of malate: X-ray crystal structure of polymeric homochiral S-malato nickel(II),  $[\Delta\text{-Ni}(\text{S-Hmal})(\text{H}_2\text{O})_2]_n \cdot n\text{H}_2\text{O}$ . Polyhedron, 2002, 21, pp. 787–790. [http://dx.doi.org/10.1016/S0277-5387\(02\)00845-8](http://dx.doi.org/10.1016/S0277-5387(02)00845-8).
- Lah, N.; Kralj, C.I.; Leban, I. Solvothermal synthesis of a novel mixed valence Cu(I)/Cu(II) complex containing sulphate, malate and 4,4'-bipyridine,  $[\text{Cu}^{\text{I}}\text{Cu}^{\text{II}}_2(\text{mal})(\text{SO}_4)(\text{bipy})_2 \cdot \text{H}_2\text{O}]_n$ . Unique binding mode of the malate anion. Inorganic Chemistry Communications, 2003, 6, pp. 1441–1444. <http://dx.doi.org/10.1016/j.inoche.2003.08.029>.
- Cunha-Silva, L.; Shi, F.N.; Paz, F.A.A.; Hardie, M.J.; Klinowski, J.; Trindade, T.; Rocha, J. Supramolecular salts containing the anionic  $[\text{Ge}(\text{C}_2\text{O}_4)_3]_2$  complex and heteroaromatic amines. Inorganica Chimica Acta, 2009, 362, pp. 263–270. <http://dx.doi.org/10.1016/j.ica.2008.02.038>.
- Shi, F.-N.; Cunha-Silva, L.; Hardie, M.J.; Trindade, T.; Paz, F.A.A.; Rocha, J. Heterodimetallic germanium(IV) complex structures with transition metals. Inorganic Chemistry, 2007, 46, pp. 6502–6515. DOI: [10.1021/ic700507j](http://dx.doi.org/10.1021/ic700507j).
- Lu, J.; Chu, D.-Q.; Yu, J.-H.; Zhang, X.; Bi, M.-H.; Xu, J.-Q. In situ ligands syntheses involving malic acid in hydrothermal condition. Inorganica Chimica Acta, 2006, 359, pp. 2495–2500. <http://dx.doi.org/10.1016/j.ica.2006.02.043>.
- Sheldrick, G.M. A short history of SHELX. Acta Crystallographica, 2008, A64, pp. 112–122. <https://doi.org/10.1107/S0108767307043930>.
- CCDC, The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK, 44 1223 336408. E-mail: deposit@ccdc.cam.ac.uk
- Bellami, L. Infrared spectra of complex molecules. IL: Moscow, 1963, 590 p. (in Russian). DOI: [10.1007/978-94-011-6520-4](http://dx.doi.org/10.1007/978-94-011-6520-4).
- Tarasevich, B.N. Infrared spectra of basic class of organic compounds. Reference materials. Moscow, 2012, 54 p. (in Russian).